

## **Genetics of the $\alpha$ -Amylase/Trypsin inhibitor family in wheat and related species**

by **R. SANCHEZ-MONGE, F. GARCIA-OLMEDO, L. GOMEZ, J. ROYO, P. CARBONERO AND G. SALCEDO\***

Departamento de Bioquímica, Escuela Técnica Superior de Ingenieros Agrónomos,  
Ciudad Universitaria, E-28040 Madrid, Spain

**Key Words:** Multi-gene family, genomic organization,  $\alpha$ -amylase inhibitors, trypsin inhibitors, wheat, barley, rye.

### **ABSTRACT**

R. Sánchez-Monge, F. García-Olmedo, L. Gómez, J. Royo, P. Carbonero and G. Salcedo. Genetics of the  $\alpha$ -Amylase/Trypsin inhibitor family in wheat and related species. *An. Aula Dei* 19 (1-2): 195-205.

Several members of an inhibitor family of heterologous  $\alpha$ -amylases and of trypsin have been isolated from wheat, barley and rye endosperms. Comparisons of the amino acid sequences of these inhibitors together with the chromosomal locations of their structural genes in the three species, indicate that they are encoded by a disperse multi-gene family that must have originated both by translocations and intrachromosomal duplications events that took place in common ancestors of wheat, barley and probably rye. Homology among some members of the inhibitor family and various proteins of maize, ragi, castor bean and others, indicate that this protein family extends to other phylogenetically distant species.

### **INTRODUCTION**

Protein inhibitors which are active against different types of hydrolytic enzymes are widely distributed in plants (for a review see García-Olmedo et al., 1987). Only a few of these inhibitors affects endogenous enzymes and may be specifically involved in plant metabolism. A majority of the plant inhibitors are active towards heterologous proteases or  $\alpha$ -amylases from various organisms. Several lines of evidence indicate that these inhibitors play a protective role against the attacks of animal predators, insects, fungi, bacteria and virus. In this context, the following observations merit special mention: a) the levels of some proteinase inhibitors are increased in

\* To whom correspondence should be addressed.

response to mechanical lesions caused by insects or by fungal infections (Ryan, 1984); b) the correlation between trypsin inhibitor levels and pests resistance in cowpea cultivars (Gatehouse et al., 1979); c) the expression of cowpea trypsin inhibitor in transgenic plants confers resistance to two of its major insect pests (Hilder et al., 1987).

Another field of practical interest concerning plant inhibitors that inhibit mammalian enzymes is their potential incidence on animal production and human nutrition, as well as their possible use in some clinical treatments (see Buonocore and Silano, 1986, and García-Olmedo et al., 1987, for reviews).

A cereal inhibitor family of heterologous  $\alpha$ -amylases and of trypsin is among those that have been more actively studied. Since Kneen and Sandstedt (1943) described the first wheat  $\alpha$ -amylase inhibitor, it has been shown that a substantial fraction of the albumins and globulins of wheat and barley endosperms is represented by a group of homologous inhibitors of  $\alpha$ -amylases and trypsin (Buonocore et al., 1977; Shewry et al., 1984; Barber et al., 1986a; Sánchez-Monge et al., 1986a,b). Some members of this protein family can be selectively extracted by chloroform/methanol mixtures and have been designated CM-proteins (García-Olmedo and Carbonero, 1970; Rodríguez-Loperena et al., 1975; Salcedo et al., 1978, 1982).

The wheat  $\alpha$ -amylase inhibitors have been classified into three classes: monomeric, dimeric and tetrameric (Deporte et al., 1976). The first type represents monomeric variants of about Mr 12 kDa, while the dimeric one can be dissociated into identical 12 kDa subunits. The chemical and inhibitory properties of these two inhibitor classes have been extensively studied (Buonocore et al., 1977; O'Connor and McGeeney, 1981; Maeda et al., 1982), and the complete amino acid sequences of one monomeric and two dimeric inhibitors have been established (Kashlan and Richardson, 1981; Maeda et al., 1983, 1985). Although the tetrameric type has been less actively investigated, Buonocore et al. (1985) have apparently isolated an inhibitor with Mr 48 kDa, which dissociates into unidentified 12-14 kDa subunits. Reconstitution experiments carried out in our laboratory (Sánchez-Monge et al., 1987; Gómez et al., 1989) have shown that the previously isolated proteins CM2, CM3 and CM16 form *T. turgidum*, and the same ones plus CM1 and CM17 from *T. aestivum* are the subunits of wheat tetrameric inhibitors.

In cultivated barley, we have recently demonstrated the existence of  $\alpha$ -amylase inhibitors corresponding to the three classes, as well as that the previously characterized proteins CMa, Cp.b and CMd are the subunits of the tetrameric inhibitor (Sánchez-Monge et al., 1986b). Also the complete sequence of a homodimeric  $\alpha$ -amylase inhibitor 'BDAI-1' has been deduced from the nucleotide sequence of its corresponded cDNA clone (Lázaro et al., 1988).

Trypsin inhibitors that belong to the same protein family of the  $\alpha$ -amylase inhibitors mentioned above have been isolated and characterized in barley (Mikola and Souliina, 1969; Odani et al., 1983; Lazaro et al., 1985; Barber et al., 1986b) and wheat (Boisen and Djurtoft, 1981). Recently the complete amino acid sequence of a trypsin inhibitor, and the N-terminal one of an  $\alpha$ -amylase inhibitor from rye endosperm have been reported (Lyons et al., 1987).

We discuss in this paper the chromosomal organization of the multi-gene family that encodes the  $\alpha$ -amylase/trypsin inhibitor family of cereal endosperms, as well as the evolutionary implications of the structural homologies among different members of this protein family.

## RESULTS AND DISCUSSION

Two-dimensional electrophoretic maps of endosperm extracts from hexaploid wheat (*T. aestivum*, genomes AABBDD) and barley (*H. vulgare*, genomes HH), containing  $\alpha$ -amylase and trypsin inhibitors are shown in Fig. 1. The fractionation by gel filtration under non-dissociating conditions of crude inhibitor extracts, allow the classification of the proteins that appeared in the two dimensional maps of Fig. 1 as subunits of tetrameric, dimeric or monomeric  $\alpha$ -amylase inhibitors, or monomeric trypsin inhibitors (Sánchez-Monge et al., 1986b, 1987; Barber et al., 1986b; Gomez et al., 1989).

Total or N-terminal amino acid sequences for different members of the inhibitor family isolated from wheat, barley and rye endosperms are presented in Fig. 2, and the percentages of homology calculated for all possible binary comparisons are summarized in Table 1. Evolutionary implications of the observed homologies are best understood if they are considered together with the chromosomal locations of the inhibitors structural genes in the three species, and will be discussed hereafter.

The distribution among chromosomes of the multi-gene family encoding this group of inhibitors has been studied through the analysis of wheat nulli-tetrasomic and ditelosomic lines (García-Olmedo and Carbonero, 1970; Aragoncillo et al., 1975; Fra-Mon et al., 1984; Sánchez-Monge et al., 1986a), wheat-barley addition lines (Salcedo et al., 1984; Hejgaard et al., 1984a) and wheat-rye addition lines (Hejgaard et al., 1984b and authors unpublished results). A summary of these observations, together with the "in vitro" activities of the corresponding proteins is presented in Table 2. It can be seen that loci encoding monomeric and dimeric  $\alpha$ -amylase inhibitors as well as the rye trypsin inhibitor and the barley trypsin inhibitor CMe, are located in chromosome groups 3 and 6. The loci for subunits of tetrameric  $\alpha$ -amylase inhibitors and the barley trypsin inhibitor CMc are in chromosomes of groups 4 and 7 of wheat, and 4 and 1 of barley. This finding further supports the proposed homology of barley chromosome 1 with chromosome group 7 of wheat (Hart et al., 1980; Powling et al., 1981). It is to be noted that in the case of wheat, which is allohexaploid, we have failed to identify triplicate genes (one in each of the three chromosomes of each chromosome group involved), which is in line with the idea of diploidization of redundant genetic information subsequent to the allopolyploid formation (García-Olmedo et al., 1978). The fact that among the genes involved the only ones associated with the A genome are those located in chromosome 4A, for which a B genome origin has been postulated (Rayburn and Gill, 1985), indicate that the distribution of this multi-gene family in wheat could be compatible with the hypothesis of a non-random diploidization (García-Olmedo, 1968; Carbonero and García-Olmedo, 1969; Galili and Feldman, 1985).

Joint consideration of the observed sequence homologies together with the inhibitory activities and the chromosomal locations of genes encoding these proteins, allows some evolutionary conclusions concerning the origin and dispersion of this multi-gene family to be drawn. There is a greater homology between a given protein from one genome and the appropriate one from a different genome than between that protein and any other encoded in the same genome. See for instance the high degree of homology (86%) between WTAI CM1 and WTAI CM2, encoded by genes located in the short arms of chromosomes 7D and 7B, respectively, and between these proteins and BTAI CMa (82-86% homology), whose genes is located in chromosome 1 of barley (equivalent to group 7 of wheat).

Barley trypsin inhibitor BTI CMc, whose gene is in the same chromosome as the  $\alpha$ -amylase inhibitor BTAI CMa, has a higher homology to this inhibitor (78%) than to the other barley trypsin inhibitor BTI CMe (45%), whose gene is located in a different chromosome.

These observations clearly indicate that the dispersion of this multi-gene family must have occurred before the evolutionary differentiation of the genomes of barley and those included in allohexaploid wheat.

The only reported  $\alpha$ -amylase inhibitor from rye, designated RAI (Lyons et al., 1987), is 70-80% homologous to proteins WTAI CM1, WTAI CM2, and BTAI CMa, which are subunits of the tetrameric  $\alpha$ -amylase inhibitors, so it is likely that RAI is also part of a tetrameric inhibitor from rye.

Proteins encoded by genes in chromosomes of group 4 of wheat and barley may be included into two groups. Homologies between WTAI CM16, WTAI CM17 and BTAI CMb (70-83%) and between WTAI CM3 and BTAI CMd (70%) are high, while the inter-group homologies are much lower (36-45%). This suggests two distinct loci associated with group 4 chromosomes whose origin may be an intrachromosomal duplication in a common ancestor of wheat and barley. In conclusion this disperse multi-gene family must have originated both by translocations and intrachromosomal duplications events that took place in common ancestors of wheat, barley and probably rye.

Although the most prominent members of this family in wheat and barley are those so far described, there is evidence for additional members of the family. Thus, Paz-Ares et al. (1986) obtained and sequenced barley cDNA clones which coded for proteins with 35-45% homology to the above-described ones. In Fig. 3, the amino acid sequence deduced from barley cDNA clone C23 is compared with that directly determined for barley trypsin inhibitor CMe. Also included in Fig. 3A are the amino acid sequences of a maize trypsin inhibitor and of a bifunctional  $\alpha$ -amylase/trypsin inhibitor from ragi (*Eleusine coracana*). The homology between the last three proteins is fairly high (50-65%), indicating that this protein family extends to other species which are more phylogenetically distant than wheat, barley and rye. It should be also pointed out that there are weak but significant homologies between these proteins and reserve globulins of some dicotyledoneous plants and with certain non repetitive domains in prolamins (Kreis et al., 1985). In Fig. 3B, the homology between the B-region of two inhibitors of this family, a 2S globulin from castor bean and a  $\gamma$ -secalin of rye is shown.

#### ACKNOWLEDGMENTS

We acknowledge the technical assistance of D. Lamonedá. This work has been supported by Comision Asesora de Investigación Científica y Técnica grant No. PB85-0193.

#### REFERENCES

- Aragoncillo, C.; M.A. Rodríguez-Loperena; P. Carbonero; F. García-Olmedo. (1975) Chromosomal control of non-gliadin proteins from the 70% ethanol extract of wheat endosperm. **Theor. Appl. Genet.** 45: 322-326.
- Barber, D.; R. Sánchez-Monge; F. García-Olmedo; G. Salcedo; E. Mendez. (1986a) Evolutionary implications of sequential homologies among members of the trypsin/ $\alpha$ -amylase inhibitor family (CM-proteins) in wheat and barley. **Biochim. Biophys. Acta** 873: 147-151.
- Barber, D.; R. Sánchez-Monge; E. Mendez; A. Lazaro; F. García-Olmedo; G. Salcedo. (1986b) New  $\alpha$ -amylase and trypsin inhibitors among the CM-proteins of barley (*Hordeum vulgare*) **Biochim. Biophys. Acta** 869: 115-118.
- Boisen, S.; R. Djurtoft. (1981) Trypsin inhibitor from wheat endosperm: purification and characterization. **Cereal Chem.** 58: 460-463.
- Buonocore, V.; M.G. De Biasi; P. Giardina; E. Pocerio; V. Silano. (1985) Purification and properties of and  $\alpha$ -amylase tetrameric inhibitor from wheat kernel. **Biochim. Biophys. Acta** 831: 40-48.
- Buonocore, V.; T. Petrucci; V. Silano. (1977) Wheat protein inhibitors of  $\alpha$ -amylase. **Phytochemistry** 16: 811-820.
- Buonocore, V.; V. Silano. (1986) Biochemical, nutritional and toxicological aspects of alpha-amylase inhibitors from plant foods. In: *Nutritional and toxicological significance of enzyme inhibitors in foods*. Friedman, M. (ed.). Plenum Publishing Corporation, pp. 483-507.
- Campos, F.A.P.; M. Richardson. (1983) The complete amino acid sequence of the bifunctional  $\alpha$ -amylase/trypsin inhibitor from seeds of ragi (Indian finger millet; *Eleusine coracana* Goertn). **FEBS Lett.** 152: 300-304.
- Carbonero, P.; F. García-Olmedo. (1969) Purothionins in *Aegilops-Triticum* spp. **Experientia** 25: 1110.
- Deponte, R.; R. Parlamenti; T. Petrucci; V. Silano; M. Tomasi. (1976) Albumin  $\alpha$ -amylase inhibitor families from wheat flour. **Cereal Chem.** 53: 805-820.
- Fra-Mon, P.; G. Salcedo; C. Aragoncillo; F. García-Olmedo. (1984) Chromosomal assignment of genes controlling salt-soluble proteins (albumins and globulins) in wheat and related species. **Theor. Appl. Genet.** 69: 167-172.

- Galili, G.; M. Feldman. (1985) Genetic control of endosperm proteins in wheat. 3. Allocation to chromosomes and differential expression of high molecular weight glutenin and gliadin genes in intravarietal substitution lines of common wheat. **Theor. Appl. Genet.** 69: 583-589.
- García-Olmedo, F. (1968) Genetics of synthesis of  $\beta$ -sitosterol esters in wheat and related species. **Nature** 220: 1144-1145.
- García-Olmedo, F.; P. Carbonero. (1970) Homoeologous protein synthesis controlled by homoeologous chromosomes in wheat. **Phytochemistry** 9: 1495-1497.
- García-Olmedo, F.; P. Carbonero; C. Aragoncillo; G. Salcedo. (1978) Loss of redundant gene expression after poliploidization in plants. **Experientia** 34: 332-333.
- García-Olmedo, F.; G. Salcedo; R. Sánchez-Monge; L. Gomez; J. Royo; P. Carbonero. (1987) Plant proteinaceous, inhibitors of proteinases and  $\alpha$ -amylases. In: *Oxford Surveys of Plant Molecular and Cell Biology*. Vol. 4. Mifflin, B. (ed.). Oxford University Press, UK, pp. 275-334.
- Gatehouse, A.M.R.; J.A. Gatehouse; P. Dobie; A.M. Kilsminster; D. Boulter. (1979) Biochemical basis of insect resistance in *Vigna unguiculata*. **J. Sci. Food Agric.** 30: 948-958.
- Gómez, L., R. Sánchez-Monge, F. García-Olmedo, G. Salcedo. (1989). Wheat tetrameric inhibitors of insect  $\alpha$ -amylase: allopolyploid heterosis at the molecular level. **Proc. Natl. Acad. Sci. USA** 86: 3242-3246.
- Hart, G.E.; A.K.M.R. Islam; K.W. Shepherd. (1980) Use of isozymes as chromosome markers in the isolation and characterization of wheat-barley chromosomal addition lines. **Genet. Res. Camb.** 36: 311-325.
- Hejggard, J.; S.E. Bjorn; G. Nielsen. (1984a) Localization to chromosomes of structural genes for the major protease inhibitors in barley grains. **Theor. Appl. Genet.** 68: 127-130.
- Hejggard, J.; S.E. Bjorn; G. Nielsen. (1984b) Rye chromosomes carrying structural genes for the major grain protease inhibitors. **Hereditas** 101: 257-259.
- Hilder, V.A.; A.M.R. Gatehouse; S.E. Sheerman; R.F. Barker; D. Boulter. (1987) A novel mechanism of insect resistance engineered into tobacco. **Nature** 300: 160-163.
- Kashlan, N.; M. Richardson. (1981) The complete amino acid sequence of a major wheat protein inhibitor of  $\alpha$ -amylase. **Phytochemistry** 20: 1781-1784.
- Kneen, E.; R.M. Sandstedt. (1943) An amylase inhibitor from certain cereals. **J. Am. Chem. Soc.** 65: 1247-1252.
- Kreis, M.; B.G. Forde; S. Rahman; B.J. Mifflin; P.R. Shewry. (1985) Molecular evolution of the seed storage proteins of barley, rye and wheat. **J. Mol. Biol.** 183: 499-502.
- Lázaro, A.; D. Barber; G. Salcedo; E. Mendez; F. García-Olmedo. (1985) Differential effects of high-lysine mutations on the accumulation of individual members of a group of proteins encoded by a disperse multigene family in the endosperm of barley (*Hordeum vulgare* L.). **Eur. J. Biochem.** 149: 617-623.
- Lázaro, A.; R. Sánchez-Monge; G. Salcedo; J. Paz-Ares; P. Carbonero; F. García-Olmedo. (1988) A dimeric inhibitor of insect  $\alpha$ -amylase from barley. Cloning of the cDNA and identification of the protein. **Eur. J. Biochem.** 172: 129-134.
- Lyons, A.; M. Richardson; S. Tatham; P.R. Shewry. (1987) Characterization of homologous inhibitors of trypsin and  $\alpha$ -amylase from seeds of rye (*Secale cereale*). **Biochim. Biophys. Acta** 916: 305-310.
- Maeda, K.; T. Hase; H. Matsubara. (1983) Complete amino acid sequence of an  $\alpha$ -amylase inhibitor in wheat kernel. **Biochim Biophys. Acta** 743: 52-57.
- Maeda, K.; S. Kakabayashi; H. Matsubara. (1985) Complete amino acid sequence of an  $\alpha$ -amylase inhibitor in wheat kernel (0.19-inhibitor). **Biochim. Biophys. Acta** 828: 213-221.
- Maeda, K.; Y. Takamori; O. Oka. (1982) Isolation and properties of an  $\alpha$ -amylase inhibitor (0.53) from wheat (*Triticum aestivum*). **Agric. Biol. Chem.** 46: 2873-2875.
- Mahoney, W.C.; M.A. Hermanson; B. Jones; D.D. Powers; R.S. Corfman; G.R. Reek. (1984) Amino acid sequence and secondary structural analysis of the corn inhibitor of trypsin and activated Hageman factor. **J. Biol. Chem.** 259: 8412-8416.

- Mikola, J.; E.M. Soulinna. (1969) Purification and properties of a trypsin inhibitor from barley. **Eur. J. Biochem.** 9: 555-560.
- O'Connor, C.M.; K.F. McGeeney. (1981) Isolation and characterization of four inhibitors from wheat flour which display differential inhibition specificities for human salivary and human pancreatic  $\alpha$ -amylases. **Biochim. Biophys. Acta** 658: 397-405.
- Odani, S.; T. Koide; T. Ono. (1983) The complete amino acid sequence of barley trypsin inhibitor. **J. Biol. Chem.** 258: 7998-8003.
- Paz-Ares, J.; F. Ponz; P. Rodríguez-Palenzuela; A. Lazaro; C. Hernández-Lucas; F. García-Olmedo; P. Carbonero. (1986) Characterization of cDNA clones of the family of trypsin/ $\alpha$ -amylase inhibitors (CM-proteins) in barley (*Hordeum vulgare* L.). **Theor. Appl. Genet.** 71: 842-846.
- Powling, A.; A.K.M.R. Islam; K.W. Shepherd. (1981) Isozymes in wheat-barley hybrid derivative lines. **Biochem. Genet.** 19: 237-254.
- Rayburn, A.L.; B.S. Gill. (1985) Molecular evidence for the origin and evolution of chromosome 4A in polyploid wheats. **Can. J. Genet. Cytol.** 27: 246-250.
- Rodríguez-Loperena, M.A.; C. Aragoncillo; P. Carbonero; F. García-Olmedo. (1975) Heterogeneity of wheat endosperm proteolipids (CM-proteins). **Phytochemistry** 14: 1219-1223.
- Ryan, C.A. (1984) Defense responses of plants. In: *Plant gene research: genes involved in microbe-plant interaction*. Verma, D.P.S.; Hohn, T.H. (eds.). Springer Verlag, Wien, pp. 375-386.
- Salcedo, G.; P. Fra-Mon; J.L. Molina-Cano; C. Aragoncillo; F. García-Olmedo. (1984) Genetics of CM-proteins (A-hordeins) in barley. **Theor. Appl. Genet.** 68: 53-59.
- Salcedo, G.; M.A. Rodríguez-Loperena; C. Aragoncillo. (1978) Relationships among low MW hydrophobic proteins from wheat endosperm. **Phytochemistry** 17: 1491-1494.
- Salcedo, G.; R. Sánchez-Monge; C. Aragoncillo. (1982) The isolation and characterization of low molecular weight hydrophobic salt-soluble proteins from barley. **J. Exp. Bot.** 33: 1325-1331.
- Sánchez-Monge, R.; D. Barber; E. Mendez; F. García-Olmedo; G. Salcedo. (1986a) Genes encoding  $\alpha$ -amylase inhibitors are located in the short arms of chromosomes 3B, 3D and 6D of wheat (*Triticum aestivum* L.). **Theor. Appl. Genet.** 72: 108-113.
- Sánchez-Monge, R.; J.A. Fernández; G. Salcedo. (1987) Subunits of tetrameric  $\alpha$ -amylase inhibitors of *Hordeum chilense* are encoded by genes located in chromosomes 4Hch and 7Hch. **Theor. Appl. Genet.** 74: 811-816.
- Sánchez-Monge, R.; L. Gomez; F. García-Olmedo; G. Salcedo. (1986b) A tetrameric inhibitor of insect  $\alpha$ -amylase from barley. **FEBS Lett.** 207: 105-109.
- Sharief, F.S.; S.S.L. Li. (1982) Amino acid sequence of small and large subunits of seed storage protein from *Ricinus communis*. **J. Biol. Chem.** 257: 14753-14759.
- Shewry, P.R.; D. Lafiandra; G. Salcedo; C. Aragoncillo; F. García-Olmedo; E.J. Lew; M.D. Dietler; D.D. Kasarda. (1984) N-terminal amino acid sequences of chloroform/methanol-soluble proteins and albumins from endosperm of wheat, barley and related species. **FEBS Lett.** 175: 359-363.

**Table 1.-** Binay comparisons (% homology) of wheat, barley and rye members of the  $\alpha$ -amylase/trypsin inhibitor family. Homology was calculated for common segments of the sequences that appear in Fig. 2. Abbreviations are as in Fig. 2.

PROTEINS	CM16	CM17	CMb	CM3	CMd	CM2	CM1	CMa	RAI	CMc	0.53	0.19	0.28	BDAI	CMe	RTI
WTAI CM16		78	83	45	40	45	41	38	38	33	27	27	27	32	38	35
WTAI CM17			70	37	36	37	37	35	29	33	26	26	30	30	37	33
BTAI CMb				52	47	45	41	38	38	33	30	30	30	29	45	38
WTAI CM3					70	50	39	43	41	38	41	41	30	26	46	48
BTAI CMd						47	37	40	45	43	30	30	23	22	43	45
WTAI CM2							86	86	80	72	38	34	37	39	54	41
WTAI CM1								82	74	69	34	31	33	35	53	45
BTAI CMa									74	78	31	28	20	32	50	41
RAI									64	25	25	29	41	41		
BTI CMc											31	28	30	32	45	25
WDAI 0.53												94	58	45	28	20
WDAI 0.19													56	44	30	17
WMAI 0.28														63	25	17
BDAI															23	17
RTI																76

Table 2.- Chromosomal locations of genes and inhibitory activities of the  $\alpha$ -amylase/trypsin inhibitors from wheat, barley and rye.

Protein grouped by loci*		Chromosome** (or chromosome arm)	Inhibitory activity	References
WDAI	0.53	3BS	$\alpha$ -amylase dimeric	a,b,c
WDAI	0.19	3DS		a,b,c
BDAI		?H		d
BTI	CMc	3H	trypsin	e,f
RTI		3R		g
WMAI	0.28	6DS	$\alpha$ -amylase monomeric	b,c
WTAI	CM16	4AS	$\alpha$ -amylase tetrameric	a,b
WTAI	CM17	4DS		a,b
BTAI	CMb	4H		f
WTAI	CM3	4AS	$\alpha$ -amylase tetrameric	a,b
WTAI	CMd	4H		f
WTAI	CM2	7BS		a,b,h
WTAI	CM1	7DS		a,b,h
BTAI	CMa	1H		f
RAI		?R		i
BTI	CMc	1H	trypsin	f

\* Abbreviations as in Fig. 2.

\*\* Hexaploid wheat (*T. aestivum*): genomes AABBDD; barley (*H. vulgare*): genomes HH, and rye (*S. cereale*): genomes RR. Barley chromosome 1 is homologous to chromosome group 7 of wheat.

References: a) Aragoncillo et al., 1975; b) Fra-Mon et al., 1984; c) Sánchez-Monge et al., 1986a; d) Lázaro et al., 1988; e) Hejgaard et al., 1984a) f) Salcedo et al., 1984; g) Hejgaard et al., 1984b; h) García-Olmedo and Carbonero, 1970; i) Lyons et al., 1987.






Figure 1.- Two-dimensional protein maps (electrofocusing, IEF, pH 4-9 x starch gel electrophoresis, SGE, pH 3.2) of crude 70% ethanol extracts from: A) hexaploid wheat (*T. aestivum* cv. Chinese Spring) and B) barley (*H. vulgare* cv. Bomi) endosperms. Proteins whose amino acid sequences appear in Fig. 2 are labeled MI, DI and TI = components of monomeric, dimeric and tetrameric  $\alpha$ -amylase inhibitors, respectively; Tr = Trypsin inhibitor. BDAI = barley dimeric  $\alpha$ -amylase inhibitor.

Inhibitors	Ref.	
WTAI CM16	1	I-GNE-DCTPWMSTLITPLPSCRDYVEEQQAC
WTAI CM17	1	NE-DCTPWTSTLIXPLPXCRNYVXXQAC
BTAI CMb	2	V-GSE-DCTPWTATPIITPLPSCRDYVEEQQAC
WTAI CM3	3	S-GS---CVPGVAFRTNLLPHCRDYVLQQTTC
BTAI CMd	3	A-AAATDCSPGVAFPTNLLGHCRDYVLQQTTC
WTAI CM2	1	T-GPY--CYPGMGLPSNPLEGCREYVAQQTTC
WTAI CM1	1	T-GPY--CYAGMGLPINPLEGCREYVASQXC
BTAI CMa	2	T-GQY--CYAGMGLPSNPLQGCREYVAZQTTC
RAI	4	T-GPY--CYAGMGLPTKPLGGCREYVAQQTTC
BTI CMc	2	T-SIYT-CYEGMGLPVNPLQGCRFYVAXQTTC
WDAI 0.53	5	S-GPWM-CYPGQAFQVPALPGCRPLL-KLQC-NG
WDAI 0.19	5	S-GPWM-CYPGQAFQVPALPACRPLL-RLQC-NG
WMAI 0.28	6	S-GPWSWCNPTATGYKSALTGCRAMV-KLQC-VG
BDAI	7	S-GPWMWCDPEMGHKVSPLTRCRALV-KLEC-VG
BTI CMe	8	F-GDS--CAPGDALPHNPLRACRTYVVSQICHQG
RTI	4	SVGGQ--CVPGLAMPHNPLGACRTYVVSQICHVG

WDAI 0.53	--SQVPEAVLRDCCQQLADTI-SEWPRCGAL-YSMLDS--MYKEHGVSEGGAGT
WDAI 0.19	--SQVPEAVLRDCCQQLAHTI-SEWCRCGAL-YSMLDS--MYKEHGAQEGGAGT
WMAI 0.28	--SQVPEAVLRDCCQQLADINNEWCRCGDL-SSMLRA--VYQELGVREGKE-V
BDAI	--NRVPEADVLRDCCQEVANISNEWCRCGDL-GSMLRS--VYAALGVGGGPEEV
BTI CMe	PRLLTSDMKRR-CCDELLSAI-PAYCRCEALRIIMO-G--VVTWQGAFFG-AYF
RTI	PRLFTWDMKRR-CCDELLSAI-PAYCRCEALRILMD-G--VVTQQGVFEG-GYL

WDAI 0.53	GAFPSCRREVVKLTAASITAV-CRL-PIVVVDASGDGAYVCKDV--AAYPDA
WDAI 0.19	GAFPRCRREVVKLTAASITAV-CRL-PIVVVDASGDGAYVCKDV--AAYPDA
WMAI 0.28	--LPGCRKEVMKLTAA-SVPEV-CKV-PIPNP-SGDRAGVCYGDW-CAYPDV
BDAI	--FPGCQKDVMMKLLVA-GVPAL-CNV-PIPNPNEAGTR-GVCY--W-SASTDT
BTI CMe	KDSPNCPREERQTSYAAANLVTPQECNLGTI--H--G-SAY-CPELQPGY-G
RTI	KDMPNCPREVTQRSYAAATLVAPQECNLPTI--H--G-SPY-CPTLQAGY

Figure 2.- Alignment of amino acid sequences of members of the  $\alpha$ -amylase/trypsin inhibitors family from wheat, barley and rye endosperm. Gaps introduced for the alignment are indicate (-). Conserved positions are boxed. W = wheat; B = barley and R = rye. TAI, DAI and MAI = subunits of tetrameric, dimeric and monomeric  $\alpha$ -amylase inhibitor, respectively. TI = trypsin inhibitor. References: (1) Barber et al., 1986b; (2) Barber et al., 1986a; (3) Shewry et al., 1984; (4) Lyons et al., 1987; (5) Maeda et al., 1985; (6) Kashlan and Richardson, 1981; (7) Lazaro et al., 1988; (8) Odani et al., 1983.

**A**

C23	LESVKDE	COLGVDF	PHNPL	AT	CHTYVIK	RV	CGRG	-	P	-	S	R	P	M	L	V	K
BTI CMe	F	GDS	-	C	A	P	G	D	A	L	P	H	N	P	L	R	A
RgBI	S	V	G	T	S	-	C	I	P	G	M	A	I	P	H	N	P
MTI	S	A	G	T	S	-	C	V	P	G	W	A	I	P	H	N	P

C23	E	R	C	C	R	E	L	A	A	M	P	D	H	C	R	C	E	A	L	R	I	L	M	D	G	-	V	R	T	P	E	G	R	V	V	E	G	R	-	L	G	D	R
BTI CMe	R	R	C	C	D	E	L	S	A	I	P	A	Y	C	R	C	E	A	L	R	I	I	M	Q	G	-	V	V	T	W	Q	G	-	A	F	E	G	A	Y	F	K	D	S
RgBI	A	R	C	C	R	Q	L	E	A	I	P	A	Y	C	R	C	E	A	V	R	I	L	M	D	G	-	V	V	T	P	S	G	-	Q	H	E	G	R	L	L	O	D	
MTI	R	R	C	C	R	E	L	A	D	I	P	A	Y	C	R	C	T	A	L	S	I	L	M	D	G	A	I	P	P	G	P	D	A	-	Q	L	E	G	-	A	L	E	

C23	D	C	P	R	E	E	Q	P	A	F	A	A	T	L	V	T	A	A	E	C	N	L	S	S	V	Q	E	P	G	V	R	-	L	V	L	A	D	G	
BTI CMe	N	C	P	R	E	R	Q	T	S	V	A	A	N	L	V	T	P	Q	E	C	N	L	G	T	I	-	H	-	G	-	S	A	Y	-	C	P	E	L	Q
RgBI	G	C	P	R	Q	V	R	A	F	A	P	K	L	V	T	E	V	E	C	N	L	A	T	I	-	H	-	G	-	G	P	F	-	C	L	S	L	-	
MTI	G	C	P	R	A	V	Q	Q	G	F	A	A	T	L	V	T	E	A	E	C	N	L	E	T	I	-	S												

**B**

BTI CMe	R	R	C	C	D	E	L	S	A	I	-	P	A	Y	C	R	C	E	A	L	R	I	I	M	Q	-	G	V	V	T	-	W	Q	G	A	F	E	G
WMAI 0.28	R	D	C	C	Q	Q	L	A	D	I	N	N	E	W	C	R	C	G	D	L	-	S	S	M	L	R	A	V	Y	Q	-	E	L	G	V	R	E	
$\gamma$ -sec.	Q	Q	C	C	Q	Q	L	A	Q	I	-	P	H	H	L	Q	C	A	A	H	S	V	V	H	-	A	I	M	Q	-	Q	E	Q	R	E	G		
2SC	R	G	C	C	D	H	L	K	Q	M	-	Q	S	Q	C	R	C	E	G	L	R	Q	A	I	Q	-	Q	Q	L	Q	G	N	M	F	E	A		

Figura 3.- Alignment of the following amino acid sequences. A) C23 = sequence deduced from barley cDNA clon pUP23 (Paz-Ares et al., 1986); BTI CMe = barley trypsin inhibitor CMe (Odani et al., 1983); RgBI = Ragi (*Eleusine coracana*) bifunctional  $\alpha$ -amylase/trypsin inhibitor (Campos and Richardson, 1983); MTI = maize trypsin inhibitor (Mahoney et al., 1984). B) B-regions as defined by Kreis et al. (1985) of: BTI CMe = barley trypsin inhibitor; WMAI 0.28 = wheat monomeric  $\alpha$ -amylase inhibitor 0.28 (Kashlan and Richardson, 1981);  $\gamma$ -sec. =  $\gamma$ -secalin from rye (Kreis et al., 1985); 2SC = 2S storage globulin from castor bean (Sharief and Li, 1982). Gaps introduced for the alignment are indicate (-). Conserved positions are boxed.